


## CHAPTER 7. THE DEFINITE INTEGRAL

### Section 7.1 Using Antiderivatives to find Area

Quick Quiz:

1. The Max-Min Theorem guarantees numbers  $m \in [x, x+h]$  and  $M \in [x, x+h]$  for which  $f(m)$  is the minimum value of  $f$  on  $[x, x+h]$ , and  $f(M)$  is the maximum value of  $f$  on  $[x, x+h]$ .
2. If  $f$  is continuous at  $a$ , then as  $x \rightarrow a$ , it must be that  $f(x) \rightarrow f(a)$ .
3. Any sketch where  $f$  IS defined at  $a$ , but  $f$  is NOT continuous at  $a$ , will work!



4.  $F(x) = x^3$  is an antiderivative of  $f(x) = 3x^2$ . Then, the desired area is given by:  $F(2) - F(0) = 2^3 - 0^3 = 8$
5. The desired area is given by:  $F(d) - F(c)$

END-OF-SECTION EXERCISES:

1.



approximation by a triangle:  $\frac{1}{2}(1)(e-1) \approx 0.86$

actual area: Using integration by parts, an antiderivative of  $f(x) = \ln x$  is  $F(x) = x \ln x - x$ . Then:

$$F(e) - F(1) = (e \ln e - e) - (1 \ln 1 - 1) = (e - e) - (0 - 1) = 1$$

3.



approximation by a trapezoid:  $\frac{1}{2}(4-1)(1+2) = \frac{1}{2}(9) = \frac{9}{2} = 4.5$

actual area: An antiderivative of  $f(x) = \sqrt{x} = x^{1/2}$  is  $F(x) = \frac{2}{3}x^{3/2} = \frac{2}{3}\sqrt{x^3}$ . Then:

$$F(4) - F(1) = \frac{2}{3}\sqrt{4^3} - \frac{2}{3}\sqrt{1^3} = \frac{2}{3}(8) - \frac{2}{3}(1) = \frac{2}{3}(7) = \frac{14}{3} \approx 4.67$$

### Section 7.2 The Definite Integral

Quick Quiz:

1. The indefinite integral  $\int f(x) dx$  gives all the antiderivatives of the function  $f$ ; by the Fundamental Theorem of Integral Calculus, if just *one* of these antiderivatives is known, then the definite integral  $\int_a^b f(x) dx$  can be computed!
2. See page 409.
3. The notation  $F(x) \Big|_a^b$  means  $F(b) - F(a)$ .
4.  $\int_{-1}^2 x^2 dx = \frac{x^3}{3} \Big|_{-1}^2 = \frac{1}{3}(2^3 - (-1)^3) = \frac{1}{3}(8 - (-1)) = \frac{1}{3}(9) = 3$
5.  $\int_{-1}^1 x^3 dx = \frac{x^4}{4} \Big|_{-1}^1 = \frac{1}{4}(1^4 - (-1)^4) = 0$ . On the interval  $[-1, 1]$ , there is the same amount of area *above* the graph of  $y = x^3$ , as there is *below*.

END-OF-SECTION EXERCISES:

1.  $\frac{48}{5}$
3.  $-6$
5.  $\frac{1}{3} \ln 2$
7.  $1 + e^2$
9. The desired area is:  $\frac{19}{24} + \frac{81}{8} = \frac{131}{12}$

### Section 7.3 The Definite Integral as the Limit of Riemann Sums

Quick Quiz:

1. A *partition* of an interval  $[a, b]$  is a finite set of points from  $[a, b]$  that includes both  $a$  and  $b$ .
2. The length of the *longest* subinterval must be  $\frac{1}{2}$ :

$$P_1 = \{1, 1.5, 2, 2.5, 3\}$$

$$P_2 = \{1, 1.3, 1.5, 2, 2.5, 3\}$$



3. There is NOT a unique Riemann sum for  $f$  corresponding to this partition; any number  $x_1^*$  may be chosen from the subinterval  $[0, 1)$ ; any number  $x_2^*$  may be chosen from the second subinterval  $[1, 2)$ , etc.
4. Think of a rectangle with 'width'  $dx$  and 'height'  $f(x)$ , where  $x$  is a number between  $a$  and  $b$ .

END-OF-SECTION EXERCISES:

1. EXP
3. SENTENCE; TRUE
5. SENTENCE; TRUE
7. SENTENCE; TRUE
9. SENTENCE; TRUE

### Section 7.4 The Substitution Technique applied to Definite Integrals

Quick Quiz:

$$1. \int (2x - 1)^3 dx = \frac{1}{2} \int u^3 du = \frac{1}{2} \frac{u^4}{4} + C = \frac{1}{8} (2x - 1)^4 + C ;$$

$$u = 2x - 1$$

$$\int_0^{1/2} (2x - 1)^3 dx = \frac{1}{8} (2x - 1)^4 \Big|_0^{1/2} = \frac{1}{8} (0 - 1) = -\frac{1}{8}$$

$$du = 2dx$$

$$2. \int_0^{1/2} (2x - 1)^3 dx = \frac{1}{2} \int_0^{1/2} (2x - 1)^3 2 dx = \frac{1}{2} \int_{-1}^0 u^3 du = \frac{1}{2} \frac{u^4}{4} \Big|_{-1}^0 = \frac{1}{8} (0 - 1) = -\frac{1}{8}$$

$$x=0 \Rightarrow u=-1$$

$$x=\frac{1}{2} \Rightarrow u=0$$

$$3. \quad u = \ln x \quad du = dx$$

$$du = \frac{1}{x} dx \quad u = x$$

$$\int_1^e \ln x dx = x \ln x \Big|_1^e - \int_1^e x \cdot \frac{1}{x} dx$$

$$= (e \ln e - 1 \ln 1) - x \Big|_1^e$$

$$= e - (e - 1) = 1$$

END-OF-SECTION EXERCISES:

1. 0
3.  $\approx 0.024$
5.  $\approx 1.931$

### Section 7.5 The Area Between Two Curves

Quick Quiz:


$$1. \int_c^d (g(x) - f(x)) dx$$

2. The  $x$ -axis is described by  $y = 0$ . The intersection points are found by:

$$-x^2 + 1 = 0 \iff x^2 = 1 \iff x = \pm 1$$

Using symmetry, the desired area is:

$$2 \int_0^1 (-x^2 + 1) dx = \left( x - \frac{x^3}{3} \right) \Big|_0^1 = 1 - \frac{1}{3} = \frac{2}{3}$$




3. A quick sketch shows that the phrase IS ambiguous; there are two regions with the indicated boundaries. Which is desired? Or, are both desired?



4.

$$\begin{aligned} \int_0^1 (e^x - (-x)) dx &= \int_0^1 (e^x + x) dx \\ &= \left( e^x + \frac{x^2}{2} \right) \Big|_0^1 \\ &= \left( e + \frac{1}{2} \right) - e^0 = e + \frac{1}{2} - 1 = e - \frac{1}{2} \end{aligned}$$




END-OF-SECTION EXERCISES:

1.  $\frac{2}{15}$
3.  $\frac{32}{3}$
5.  $\approx 2.438$
7.  $20\frac{1}{4}$



Section 7.6 Finding the Volume of a Solid of Revolution—Disks

Quick Quiz:


1. Revolve  $x = r$  about the  $y$ -axis; or revolve  $y = r$  about the  $x$ -axis.



2. Revolve  $y = -\frac{h}{r}x + h$  about the  $y$ -axis; or revolve  $y = \frac{h}{r}x$  about the  $y$ -axis. (There are other correct answers.)



3.




$$\int_0^1 \pi(x^2)^2 dx = \pi \frac{x^5}{5} \Big|_0^1 = \frac{\pi}{5}(1 - 0) = \frac{\pi}{5}$$

4. intersection points of  $y = x^2$  and  $y = 1$ :  $x^2 = 1 \iff x = \pm 1$

Also:  $y = x^2 \iff x = \pm\sqrt{y}$

A typical 'slice' at a distance  $y$  has volume  $\pi(\sqrt{y})^2 dy$ . The desired volume is:

$$\int_0^1 \pi(\sqrt{y})^2 dy = \int_0^1 \pi y dy = \pi \frac{y^2}{2} \Big|_0^1 = \frac{\pi}{2}(1 - 0) = \frac{\pi}{2}$$



## END-OF-SECTION EXERCISES:


1.  $\frac{4\pi}{3}$
3.  $\frac{\pi}{2}$
5.  $8\pi$
7.  $\frac{128\pi}{5}$
9.  $\frac{8\pi}{3}$
11.  $\frac{\pi}{4}$

## Section 7.7 Finding the Volume of a Solid of Revolution—Shells

Quick Quiz:

1. 'Cut' the shell and unroll it; the volume is:

$$2\pi r \cdot h \cdot dx$$



2.

$$\int_0^2 2\pi x(x) dx = 2\pi \frac{x^3}{3} \Big|_0^2 = \frac{2\pi}{3} (8 - 0) = \frac{16\pi}{3}$$

3. To use horizontal disks would require disks 'with holes'. Thus, in this case, shells are easier to use.

## END-OF-SECTION EXERCISES:

1.  $\frac{4\pi}{3}$
3.  $2\pi$