





SECTION 5.3 The Second Derivative—Inflection Points

IN-SECTION EXERCISES:

EXERCISE 1.



EXERCISE 2.



EXERCISE 3.

- If $x = 2$, then $x^2 = 4$; true
Converse: If $x^2 = 4$, then $x = 2$; false. (Take $x = -2$. Then the hypothesis ' $(-2)^2 = 4$ ' is true, but the conclusion ' $-2 = 2$ ' is false.)
- $1 = 2 \implies 1 + 1 = 2$; (vacuously) true
Converse: $1 + 1 = 2 \implies 1 = 2$; false
- If $1 = 2$, then $2 = 3$; (vacuously) true
Converse: If $2 = 3$, then $1 = 2$; (vacuously) true
- $A \implies B$ has converse $B \implies A$ which has converse $A \implies B$. Thus, the converse of the converse is the original implication.
original implication: $A \implies B$;
converse: $B \implies A$;
contrapositive of the converse: $\text{not } A \implies \text{not } B$
original implication: $A \implies B$;
contrapositive: $\text{not } B \implies \text{not } A$;
converse of the contrapositive: $\text{not } A \implies \text{not } B$

EXERCISE 4.

- $f'(x) = 4x$; $f''(x) = 4$. When x changes by an amount Δx , the slopes of the tangent lines should change by four times this amount.
- At $x + \Delta x$, the slope of the tangent line is $f'(x + \Delta x) = 4(x + \Delta x) = 4x + 4\Delta x$. At x , the slope of the tangent line is $f'(x)$.
-

$$\begin{aligned} \Delta f' &= f'(x + \Delta x) - f'(x) \\ &= (4x + 4\Delta x) - 4x \\ &= 4\Delta x \end{aligned}$$

Thus, the slopes of the tangent lines have indeed changed by four times the amount that x has changed.

EXERCISE 5.

- $f'(x) = 3x^2$; $f''(x) = 6x$. At the point $(2, 8)$, the slopes of the tangent lines are changing $f''(2) = 6 \cdot 2 = 12$ times as fast as x changes.
- In moving from $x = 2$ to $x = 2.1$, the change in x is 0.1 ; thus, the expected change in the slopes is: $12 \cdot 0.1 = 1.2$


3. $f'(2) = 3(2)^2 = 12$; $f'(2.1) = 3(2.1)^2 = 13.23$

Thus: $\Delta f' = 13.23 - 12 = 1.23$

4. The estimate was a bit low. This is because, as soon as we move away from the point $(2, 8)$, the rate of change of the slopes is actually *greater than* 12.

EXERCISE 6.

1. Start with $y = x^4$; shift 4 to the right; make all the y -values negative; shift up 20



2. $f(2) = -(2 - 4)^4 + 20 = 4$

3. $f'(x) = -4(x - 4)^3$; $f''(x) = -12(x - 4)^2$

Thus: $f''(2) = -12(2 - 4)^2 = -48$

When x changes by a small amount, we expect the slopes of the tangent lines to change by -48 times this amount.

4. When x changes by 0.1, the slopes should change by approximately: $-48 \cdot (0.1) = -4.8$

5. $f'(2) = -4(2 - 4)^3 = 32$; $f'(2.1) = -4(2.1 - 4)^3 = 27.436$

Note that:

$$f'(2.1) - f'(2) = 27.436 - 32 = -4.564$$

Compare this with the 'expected' change of -4.8 .

EXERCISE 7.

1. The domain of P is \mathbb{R} .

$$P'(x) = 4x^3 - 12x^2 - 7$$

$$P''(x) = 12x^2 - 24x = 12x(x - 2)$$


$$P''(x) = 0 \iff x = 0 \text{ or } x = 2$$

When $x = 0$, $P(0) = 1$, so $(0, 1)$ is a candidate for an inflection point.

When $x = 2$, $P(2) = -29$, so $(2, -29)$ is a candidate for an inflection point.

Determine the sign of P'' everywhere:

Test Points: $P''(-1) = (-)(-) > 0$, $P''(1) = (+)(-) < 0$, $P''(3) = (+)(+) > 0$



Thus, both candidates are indeed inflection points, since the concavity of the function *changes* as we pass through each point.

2. The domain of f is $(0, \infty)$.

$$\begin{aligned} f'(x) &= \frac{1}{2}x^{-1/2} + 2x \\ f''(x) &= -\frac{1}{4}x^{-3/2} + 2 \\ &= -\frac{1}{4\sqrt{x^3}} + 2 \cdot \frac{4\sqrt{x^3}}{4\sqrt{x^3}} \\ &= \frac{-1 + 8\sqrt{x^3}}{4\sqrt{x^3}} \end{aligned}$$


Note that:

$$f''(x) = 0 \iff -1 + 8\sqrt{x^3} = 0 \iff \sqrt{x^3} = \frac{1}{8} \iff x^3 = \frac{1}{64} \iff x = \frac{1}{4}$$

The point $(\frac{1}{4}, f(\frac{1}{4})) = (\frac{1}{4}, \frac{9}{16})$ is the only candidate for an inflection point.

Determine the sign of f'' everywhere:

Test Points: $f''(\frac{1}{8}) < 0$, $f''(1) > 0$



Thus, $(\frac{1}{4}, \frac{9}{16})$ is indeed an inflection point.

EXERCISE 8.

Some approximation is necessary. It is assumed that the patterns displayed at the graph boundaries continue.

- The function is positive on the interval $(-1.75, \infty)$
The function is negative on $(-\infty, -1.75)$
- The function increases on $(-\infty, 1) \cup (2.7, \infty)$
The function decreases on $(1, 2.7)$
- It is somewhat difficult to tell where the concavity changes, without further information. Thus, the following answers are certainly approximate:
The function is concave up on $(-1, 0) \cup (2, \infty)$
The function is concave down on $(-\infty, -1) \cup (0, 2)$

EXERCISE 9.

- Proof.** Suppose that $f'(c) = 0$ and $f''(c) < 0$. Assume, for simplicity, that f is defined on both sides of c .

Recall that $f'' = (f')'$. Thus, $f''(c) < 0$ means that the limit

$$\lim_{h \rightarrow 0} \frac{f'(c+h) - f'(c)}{h} = \lim_{h \rightarrow 0} \frac{f'(c+h) - 0}{h}$$

exists, and is negative. Call the value of this limit N (for 'negative'). Thus, it is possible to get the values $\frac{f'(c+h)}{h}$ as close to N as desired, merely by requiring that h be sufficiently close to 0. Remember that when h is close to 0, $c+h$ is close to c . In particular, when $h < 0$, $c+h$ is to the left of c ; and when $h > 0$, $c+h$ is to the right of c .

Refer to the sketch. Choose ϵ so that every number in the interval $I := (N - \epsilon, N + \epsilon)$ is negative. Then, find δ so that whenever h is within δ of 0, the numbers $\frac{f'(c+h)}{h}$ end up in I .

If $h < 0$, and within δ of 0, then multiplying both sides of the inequality

$$\frac{f'(c+h)}{h} < 0$$

by the negative number h yields

$$f'(c+h) > 0,$$


so the function is increasing to the left of the point $(c, f(c))$.

Similarly, if $h > 0$ and within δ of 0, then we get

$$f'(c+h) < 0,$$

so the function is decreasing to the right of the point $(c, f(c))$.

By the First Derivative Test, the point $(c, f(c))$ is a local maximum. ■



2.

$$P'(x) = 12x^3 + 12x^2 - 24x = 12x(x^2 + x - 2) = 12x(x+2)(x-1)$$

$$P''(x) = 36x^2 + 24x - 24 = 12(3x^2 + 2x - 2)$$

Now,

$$P'(x) = 0 \iff x = 0 \text{ or } x = -2 \text{ or } x = 1,$$

so there is a horizontal tangent line at each of these points.

$P''(0) = 12(0+0-2) < 0$, so P is concave down at $x = 0$. Thus, $(0, P(0))$ is a local maximum.

$P''(-2) = 12(3(-2)^2 + 2(-2) - 2) = 12(12 - 4 - 2) > 0$, so P is concave up at $x = -2$. Thus, $(-2, P(-2))$ is a local minimum.

$P''(1) = 12(3+2-2) > 0$, so P is also concave up at $x = 1$. Thus, $(1, P(1))$ is a local minimum.

END-OF-SECTION EXERCISES:

1.


$$P'(x) = 4x^3 - 6x^2 + 2x = 2x(2x^2 - 3x + 1) = 2x(2x-1)(x-1)$$

$$P''(x) = 12x^2 - 12x + 2 = 2(6x^2 - 6x + 1)$$

The critical points are summarized in the table below.

c	$P(c)$	WHY?	LOCAL EXT.?
0	10	$P'(c) = 0$	MIN
1/2	161/16	$P'(c) = 0$	MAX
1	10	$P'(c) = 0$	MIN

First Derivative Test: The sign of $P'(x)$ is given below:



Since P decreases to the left of $x = 0$ and increases to the right, there is a local minimum at $x = 0$.
 Since P increases to the left of $x = \frac{1}{2}$ and decreases to the right, there is a local maximum at $x = \frac{1}{2}$.
 Since P decreases to the left of $x = 1$ and increases to the right, there is a local minimum at $x = 1$.

Second Derivative Test: $P''(0) > 0$, so there is a local minimum at $x = 0$

$P''(\frac{1}{2}) = 2(6(\frac{1}{2})^2 - 6(\frac{1}{2}) + 1) < 0$, so there is a local maximum at $x = \frac{1}{2}$

$P''(1) > 0$, so there is a local minimum at $x = 1$


2.
$$P'(x) = 36x^3 + 48x^2 + 12x = 12x(3x^2 + 4x + 1) = 12x(3x + 1)(x + 1)$$

$$P''(x) = 108x^2 + 96x + 12$$

The critical points are summarized in the table below.

C	P(c)	WHY?	LOCAL EXT
0	1	$P'(c) = 0$	
$-\frac{1}{3}$	$\frac{32}{27}$	$P'(c) = 0$	
-1	0	$P'(c) = 0$	

First Derivative Test: The sign of $P'(x)$ is given below:



Since P decreases to the left of $x = -1$ and increases to the right, there is a local minimum at $x = -1$.
 Since P increases to the left of $x = -\frac{1}{3}$ and decreases to the right, there is a local maximum at $x = -\frac{1}{3}$.
 Since P decreases to the left of $x = 0$ and increases to the right, there is a local minimum at $x = 0$.

Second Derivative Test: $P''(-1) > 0$, so there is a local minimum at $x = -1$

$P''(-\frac{1}{3}) < 0$, so there is a local maximum at $x = -\frac{1}{3}$

$P''(0) > 0$, so there is a local minimum at $x = 0$

- $f(x)$ is positive on $(-\infty, -2.5) \cup (-2, \infty)$
 $f(x)$ is negative on $(-2.5, -2)$
- f increases on $(-4, -3) \cup (0, \infty)$
 f decreases on $(-3, -2) \cup (-2, 0)$
- f is concave up on $(-2, 2)$
 f is concave down on $(-3, -2) \cup (2, \infty)$
- $\mathcal{D}(f) = \mathbb{R} - \{-2\} = (-\infty, -2) \cup (-2, \infty)$
- $\mathcal{D}(f') = \mathbb{R} - \{-4, -3, -2\}$
- $\{x \mid f'(x) = 0\} = (-\infty, -4) \cup \{0\}$
- $\{x \mid f(x) > 10\} = (-2, -1.5)$
- $\{x \mid f'(x) > 0\} = (-4, -3) \cup (0, \infty)$
- $\{x \mid f''(x) < 0\} = (-3, -2) \cup (2, \infty)$
- $\lim_{x \rightarrow 0} f(x) = 2$
- $\lim_{t \rightarrow -2} f(t)$ does not exist
- $\lim_{y \rightarrow -4} f(y) = 4$
- The critical points are:
 $\{(x, 4) \mid x \in (-\infty, -4)\}$, since $f'(x) = 0$ for all these points
 $(0, 2)$, since $f'(0) = 0$
 $(-4, 4)$ and $(-3, 8)$, since the derivative does not exist at these points

16. $f(0) = 2$

$f'(0) = 0$

$f(1000) \approx 10$

$f'(1000) \approx 0$

17. $\{x \in \mathcal{D}(f) \mid f \text{ is not differentiable at } x\} = \{-4, -3\}$

18. The only inflection point is $(2, 6)$.

19. $\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = f'(0) = 0$

20. $\lim_{x \rightarrow -3.5} f'(x)$ equals the slope of the tangent line at $x = -3.5$ Using the points $(-4, 4)$ and $(-3, 8)$ to compute this slope, we get:

$$\frac{8 - 4}{1} = 4$$