

SECTION 7.2 The Definite Integral

IN-SECTION EXERCISES:

EXERCISE 1.

1. FALSE. The definite integral is a NUMBER.
2. TRUE
3. FALSE. The indefinite integral is a class of functions; the definite integral is a NUMBER.
4. TRUE. The actual definition is presented in section 7.3.

EXERCISE 2.

1.
$$\int_0^1 x^5 dx = \frac{1}{6}x^6 \Big|_0^1 = \frac{1}{6}(1^6 - 0^6) = \frac{1}{6}$$

Since x^5 is nonnegative on the interval $[0, 1]$, the number $\frac{1}{6}$ gives the area under the graph of $y = x^5$ on $[0, 1]$.

2.
$$\int_0^4 e^x dx = e^x \Big|_0^4 = e^4 - e^0 = e^4 - 1 \approx 53.6$$

Since e^x is positive (everywhere), the number $e^4 - 1$ gives the area under the graph of $y = e^x$ on $[0, 4]$.

3.
$$\int_{-4}^0 e^x dx = e^x \Big|_{-4}^0 = e^0 - e^{-4} = 1 - \frac{1}{e^4} \approx 0.98$$

Since e^x is positive (everywhere), the number $1 - \frac{1}{e^4}$ gives the area under the graph of $y = e^x$ on $[-4, 0]$.

4.
$$\int_1^2 \frac{1}{x} dx = \ln|x| \Big|_1^2 = \ln 2 - \ln 1 = \ln 2 \approx 0.69$$

Since $\frac{1}{x}$ is positive on $[1, 2]$, the number $\ln 2$ gives the area under the graph of $y = \frac{1}{x}$ on $[1, 2]$.

5.
$$\int_{1/2}^2 \frac{1}{x} dx = \ln|x| \Big|_{1/2}^2 = \ln 2 - \ln(1/2) = \ln 2 - \ln 2^{-1} = \ln 2 + \ln 2 = 2 \ln 2 \approx 1.39$$

Since $\frac{1}{x}$ is positive on $[\frac{1}{2}, 2]$, the number $2 \ln 2$ gives the area under the graph of $y = \frac{1}{x}$ on $[\frac{1}{2}, 2]$.

EXERCISE 3.

$\int_a^b f(x) dx = \int_a^b f(t) dt = \int_1^b f(s) ds$; only different dummy variables have been used

However, the integral $\int_a^b f(x) dx$ gives no information about $\int_c^d f(x) dx$; unless $a = c$ and $b = d$. Here, the function f is being integrated over a *different* interval.

The integral $\int_a^b f(x) dx$ gives no information about $\int_a^b g(x) dx$, unless $f = g$ on $[a, b]$. A different function is being integrated.

EXERCISE 4.

1. $\int_{-3}^{-2} f(x) dx = A$
2. $\int_{-3}^0 f(t) dt = A - 2A = -A$. The area beneath the x -axis is treated as negative by the definite integral.
3. $\int_{-3}^2 f(s) ds = A - 2A + 3A = 2A$
4. This integral cannot be computed exactly with only the given information; however, one would suspect that $\int_0^5 f(x) dx$ is a negative number.
5. $\int_{-2}^2 f(t) dt = -2A + 3A = A$
6. This integral cannot be computed exactly with only the given information; however, one would suspect that: $\int_{-3}^{-1} f(y) dy \approx 0$

EXERCISE 5.

Method I: First solve the companion indefinite integral problem:

$$\begin{aligned} u &= 5x+1 \\ du &= 5dx \end{aligned} \quad \int \frac{1}{5x+1} dx = \frac{1}{5} \int \frac{1}{u} du = \frac{1}{5} \ln |5x+1| + C$$

Then use the simplest antiderivative to evaluate the definite integral:

$$\int_0^1 \frac{1}{5x+1} dx = \frac{1}{5} \ln |5x+1| \Big|_0^1 = \frac{1}{5} (\ln 6 - \ln 1) = \frac{1}{5} \ln 6$$

Method II: Solve the definite integral directly:

$$\begin{aligned} \int_0^1 \frac{1}{5x+1} dx &= \int_0^1 \frac{1}{5(x+\frac{1}{5})} dx = \frac{1}{5} \int_0^1 \frac{1}{x+\frac{1}{5}} dx \\ &= \frac{1}{5} \ln |x+\frac{1}{5}| \Big|_0^1 = \frac{1}{5} (\ln \frac{6}{5} - \ln \frac{1}{5}) \\ &= \frac{1}{5} (\ln 6 - \ln 5 + \ln 5) = \frac{1}{5} \ln 6 \end{aligned}$$

Of course, the answers agree!

EXERCISE 6.

$$\int_0^3 (-(x-2)^2 + 1) dx = -\frac{1}{3}(x-2)^3 + x \Big|_0^3 = [-\frac{1}{3} + 3] - [-\frac{1}{3}(-8)] = \frac{8}{3} - \frac{8}{3} = 0$$

The integral is 0, because the magnitude of area above the x -axis is the same as the magnitude of the area below the x -axis, on the interval $[0, 3]$.


EXERCISE 7.

The desired area must be found in two pieces:

$$\begin{aligned} \int_{-3}^{-1} [(x+2)^2 - 1] dx &= \left[\frac{(x+2)^3}{3} - x \right] \Big|_{-3}^{-1} \\ &= \left[\frac{(-1+2)^3}{3} - (-1) \right] - \left[\frac{(-3+2)^3}{3} - (-3) \right] \\ &= \left(\frac{1}{3} + 1 \right) - \left(-\frac{1}{3} + 3 \right) = \frac{4}{3} - \frac{8}{3} = -\frac{4}{3} \end{aligned}$$

The answer is negative, since the area is beneath the x -axis.

$$\begin{aligned} \int_{-1}^0 [(x+2)^2 - 1] dx &= \left[\frac{(x+2)^3}{3} - x \right] \Big|_{-1}^0 \\ &= \frac{2^3}{3} - \left(\frac{1}{3} - (-1) \right) \\ &= \frac{8}{3} - \frac{4}{3} = \frac{4}{3} \end{aligned}$$



The desired area is: $\frac{4}{3} + \frac{4}{3} = \frac{8}{3}$

END-OF-SECTION EXERCISES:

- $\int_0^2 \frac{3}{2} x^4 dx = \frac{3}{2} \frac{x^5}{5} \Big|_0^2 = \frac{3}{10} (2^5 - 0^5) = \frac{48}{5}$
- $\int_1^8 t^{1/3} dt = \frac{3}{4} t^{4/3} \Big|_1^8 = \frac{3}{4} (8^{4/3} - 1^{4/3}) = \frac{3}{4} ((8^{1/3})^4 - 1) = \frac{3}{4} (16 - 1) = \frac{45}{4}$
- $\int_{-1}^1 (2x - 3) dx = (2 \frac{x^2}{2} - 3x) \Big|_{-1}^1 = (1 - 3) - (1 + 3) = -2 - 4 = -6$

4.
$$\int_0^1 (ax + b) dx = \left(a\frac{x^2}{2} + bx \right) \Big|_0^1 = \frac{a}{2} + b$$

5. First, find the companion indefinite integral:

$$\int \frac{x^2}{1+x^3} dx = \frac{1}{3} \int \frac{1}{1+x^3} (3x^2) dx = \frac{1}{3} \int \frac{1}{u} du = \frac{1}{3} \ln |u| + C = \frac{1}{3} \ln |1+x^3| + C$$

Now, use the simplest antiderivative to evaluate the definite integral:

$$\int_0^1 \frac{x^2}{1+x^3} dx = \frac{1}{3} \ln |1+x^3| \Big|_0^1 = \frac{1}{3} (\ln 2 - \ln 1) = \frac{1}{3} \ln 2$$

6.

$$\begin{aligned} \int_{\ln 2}^{\ln 3} e^{2t} dt &= \frac{1}{2} e^{2t} \Big|_{\ln 2}^{\ln 3} \\ &= \frac{1}{2} (e^{2 \ln 3} - e^{2 \ln 2}) = \frac{1}{2} (e^{\ln 3^2} - e^{\ln 2^2}) \\ &= \frac{1}{2} (9 - 4) = \frac{5}{2} \end{aligned}$$


7.

$$\int_0^2 (1 + e^x) dx = (x + e^x) \Big|_0^2 = (2 + e^2) - (0 + e^0) = 1 + e^2$$

8. The graph of f is a parabola that crosses the x -axis at $x = 1$ and $x = -3$. Using calculus to find the vertex: $f(x) = x^2 + 2x - 3$, so $f'(x) = 2x + 2$

$$f'(x) = 0 \iff 2x + 2 = 0 \iff x = -1$$

Also, $f(-1) = (-1 - 1)(-1 + 3) = (-2)(2) = -4$. Since $f''(x) = 2$, the graph is always concave up.



The desired area must be found in two pieces:

$$\begin{aligned} \int_{-2}^1 f(x) dx &= \int_{-2}^1 (x^2 + 2x - 3) dx = \left(\frac{x^3}{3} + x^2 - 3x \right) \Big|_{-2}^1 \\ &= \left(\frac{1}{3} + 1 - 3 \right) - \left(-\frac{8}{3} + 4 + 6 \right) \\ &= -\frac{5}{3} - \frac{22}{3} = -\frac{27}{3} = -9 \end{aligned}$$

The answer is negative, because the area lies beneath the x -axis. Then:

$$\begin{aligned} \int_1^2 (x^2 + 2x - 3) dx &= \left(\frac{x^3}{3} + x^2 - 3x \right) \Big|_1^2 = \left(\frac{8}{3} + 4 - 6 \right) - \left(\frac{1}{3} + 1 - 3 \right) \\ &= \frac{2}{3} - \left(-\frac{5}{3} \right) = \frac{7}{3} \end{aligned}$$

The desired area is: $9 + \frac{7}{3} = 11\frac{1}{3}$

9. First, factor f : find A and B with $AB = (2)(-3) = -6$ and $A + B = 5$

Take: $A = 6$ and $B = -1$

Then:

$$2x^2 + 5x - 3 = 2x^2 + 6x - x - 3 = 2x(x + 3) - (x + 3) = (2x - 1)(x + 3)$$

The graph of f is a parabola that crosses the x -axis at $x = -3$ and $x = \frac{1}{2}$.

$$f'(x) = 4x + 5$$

$$f'(x) = 0 \iff 4x + 5 = 0 \iff x = -\frac{5}{4}$$

$$f(-\frac{5}{4}) = 2(-\frac{5}{4})^2 + 5(-\frac{5}{4}) - 3 = \dots = -\frac{49}{8}$$

Since $f''(x) = 4 > 0$, the graph is concave up everywhere.

The desired area must be found in two pieces:

$$\begin{aligned} \int_0^{1/2} (2x^2 + 5x - 3) dx &= (2\frac{x^3}{3} + 5\frac{x^2}{2} - 3x) \Big|_0^{1/2} \\ &= [\frac{2}{3}(\frac{1}{2})^3 + \frac{5}{2}(\frac{1}{2})^2 - 3(\frac{1}{2})] - 0 \\ &= \frac{1}{12} + \frac{5}{8} - \frac{3}{2} = \frac{2}{24} + \frac{15}{24} - \frac{36}{24} = -\frac{19}{24} \end{aligned}$$

The answer is negative, since the area lies beneath the x -axis. Then:

$$\begin{aligned} \int_{1/2}^2 (2x^2 + 5x - 3) dx &= (2\frac{x^3}{3} + 5\frac{x^2}{2} - 3x) \Big|_{1/2}^2 \\ &= [\frac{2}{3}(2^3) + \frac{5}{2}(2^2) - 3(2)] - [\frac{2}{3}(\frac{1}{2})^3 + \frac{5}{2}(\frac{1}{2})^2 - 3(\frac{1}{2})] \\ &= \dots = \frac{81}{8} \end{aligned}$$

The desired area is: $\frac{19}{24} + \frac{81}{8} = \frac{131}{12}$

